U.S. DEPARTMENT OF THE INTERIOR
U.S. GEOLOGICAL SURVEY



Base from U.S. Geological Survey Arkansas State base map, 1990
Lambert conformal conic projection, standard parallels 33° and 45°

INTERIOR-GEOLOGICAL SURVEY, RESTON, VA 1993

MISSISSIPPI EMBAYMENT

Alluvium—Both units are equivalent in age

Qcm — **Alluvial deposits in major stream channels or in mappable meanders of major streams**—Includes alluvial deposits in natural levees in some areas

Qao — **Alluvial deposits of local streams or of overbank flow of major streams**—In some areas includes deposits in abandoned meanders of major streams

Qt — **Terrace deposits**—Alluvial deposits on one or more terrace levels

Qds — **Dune sand**—Deposited mostly on terrace deposits of the second level. Younger than some terrace deposits, older than others

Qss — **Silt and sand**—Contains lenses of gravel and clay

Ql — **Loess**

Qsg — **Sand and gravel**

Tj — **Jackson Group**

Tc — **Claiborne Group**

Tw — **Wilcox Group**

Tm — **Midway Group**

GULF COASTAL PLAIN

Qal — **Alluvium**—Alluvial deposits of present streams

Qt — **Terrace deposits**—Alluvial deposits on one or more terrace levels

Tj — **Jackson Group**

Tc — **Claiborne Group**

Tw — **Wilcox Group**

Tm — **Midway Group**

Kad — **Arkadelphia Marl**

Kn — **Nacatoch Sand**

Ks — **Saratoga Chalk**

Kmn — **Marlbrook Marl**

Ka — **Annona Chalk**

Ko — **Ozan Formation**

Kb — **Brownstown Marl**

Ktn — **Tokio Formation**

Kw — **Woodbine Formation**

Kk — **Kiamichi Formation and Goodland Limestone**

Kt — **Trinity Group (Kt)**

Kde — **Kde, De Queen Limestone**

Kdi — **Kdi, Dierks Limestone**

GULF COASTAL PLAIN, OZARK REGION, AND ARKANSAS VALLEY AND OUACHITA MOUNTAINS REGIONS

Igneous rocks—Includes undifferentiated altered sedimentary rocks of Paleozoic age at Granite Mountain, Pulaski County, and Murfreesboro, Pike County

Acid to intermediate—Nepheline syenite complexes at Magnet Cove, Hot Spring County, and Potash Sulphur Springs, Garland County, stocks at Granite Mountain, Pulaski County and in Saline County, explosion breccias, Saline and Conway Counties, and dikes and sills (principally trachyte, trequatte, phonolite, some ranging from quartz syenite to dolinite)

Basic to ultrabasic—Lamprorite pipes (Murfreesboro, Pike County) and dikes and sills (principally lamprophyres of fourchite, ouachitite, and monchiquite)

OUACHITA MOUNTAINS REGION

Ollolollths of Metagabbro and Soapstone-Serpentine—Altered peridotite

SYMBOLS

Contact

Normal Fault

Thrust fault—Also cross fault in some areas, dashed under Lake Ouachita to show continuity

OZARK REGION

Qat — **Alluvium and terrace deposits**—Alluvial deposits of present streams and on one or more terrace levels

Tg — **Gravel**—Gravel on isolated hills within 40 miles (64 km) of the Mississippi Embayment. Many small deposits to the west not shown

Kac — **Sand and clay**

Kc — **Cretaceous rocks**

Pa — **Atoka Formation**

Pbh — **Bloyd Shale, and Prairie Grove Member of the Hale Formation**

Phc — **Cane Hill Member of the Hale Formation**

Mpfb — **Pitkin Limestone, Fayetteville Shale (including the Wedington Sandstone Member), and Batesville Sandstone (including the Hindsville Limestone Member)**

Mr — **Ruddell Shale**

Mm — **Moorefield Formation**

Mb — **Boone Formation**

MDog — **Chattanooga Shale (Lower Mississippian and Upper Devonian), Clifty Limestone (Middle Devonian), and Penters Chert (Lower Devonian)**

Sslb — **Lafferty, St. Clair, and Brassfield Limestones**

Ocj — **Cason Shale and Fernvale Limestone (Upper Ordovician) and Kimmswick Limestone, Plattin Limestone, and Joachim Dolomite (Middle Ordovician)**

Ose — **St. Peter Sandstone and Everton Formation (Middle Ordovician)**

Op — **Powell Dolomite**

Oojc — **Cotter and Jefferson City Dolomites**

ARKANSAS VALLEY AND OUACHITA MOUNTAINS REGIONS

Qal — **Alluvium**—Alluvial deposits of present streams

Qt — **Terrace deposits**—Alluvial deposits on one or more terrace levels

Pbg — **Boggy Formation**

Psv — **Savanna Formation**

Pma — **McAlester Formation**

Phs — **Hartshorne Sandstone**

Pa — **Atoka Formation**, undivided

Pau — Upper part

Pam — Middle part

Pal — Lower part

Pjv — **Johns Valley Shale**

Pj — **Jackfork Sandstone**

Ms — **Stanley Shale**—Includes Chickasaw Creek Chert equivalent of Hatton (1938) near top and Hatton Tuff Lentil and Hot Springs Sandstone Member near base

MDa — **Arkansas Novaculite**—Includes upper division (Mississippian), middle division (Mississippian and Devonian), and lower division (Devonian)

Smb — **Missouri Mountain Shale and Blaylock Sandstone**—The Blaylock Sandstone is present only in the Cross, Cossatot, and Trap Mountains. Missouri Mountain Shale is mapped with Polk Creek Shale and Bigfork Chert in the area between Paron, Saline County, and Little Rock

Opb — **Polk Creek Shale (Upper Ordovician) and Bigfork Chert (Upper and Middle Ordovician)**

Ow — **Womble Shale (Middle Ordovician)**

Ob — **Blakely Sandstone (Middle and Lower Ordovician)**

Om — **Mazarn Shale**

Ocm — **Crystal Mountain Sandstone**

Oc — **Collier Shale**



MAP SHOWING SOURCES OF GEOLOGIC DATA

COMPILERS

Bush, W.V.[1]
Bush, W.V., and Haley, B.R.[2]
Clardy, B.F.[1], and Bush, W.V.
Glick, E.E.[2]
Haley, B.R.
Haley, B.R., and Zachry, D.L.[1]
Stone, C.G.[1]
Stone, C.G., and Haley, B.R.
Woodward, M.B.[2], Stone, C.G., and Haley, B.R.

[1]Arkansas Geological Commission
[2]United States Geological Survey

18 — Sources of data

Region used in Geologic Map Explanation

COMPILERS AND SOURCES

This map, like every State geologic map, is a graphic summary of information about rock units in the State at the time of the map compilation. The scale of the map forces an exaggeration of the outcrop width of the thinnest map units; therefore, rock units older than shown may be present in narrow valleys and younger than shown may be present in small outliers. The differences in tectonic features, lithologic boundaries, geographic extent of rock units, and stratigraphic nomenclature between the Geologic Map of Arkansas (Arkansas Geological Survey, 1929) and this map are based on published and unpublished geologic reports prepared since 1929 and on detailed and reconnaissance mapping done in conjunction with the preparation of this map.

Areas of responsibility for the compilation of the map are shown by color on the adjacent map. Sources cited in the list, keyed by number to the map, have been locally modified or adjusted by the compilers who also contributed detailed or reconnaissance mapping for the areas for which they are responsible.

SOURCES OF DATA

1. Adams, G.I., and Ulrich, E.O., 1905, Description of the Fayetteville quadrangle [Arkansas—Missouri]: U.S. Geol. Survey Geol. Atlas, Folio 119.
2. Arkansas Geological Commission, unpublished maps of Pointers Bluff and Guion of the White River high calcium district; Izard, Stone, and Independence Counties, Arkansas, 1958–59.
3. Arndt, H.H., U.S. Geol. Survey, unpublished data.
4. Dane, C.H., 1929, Upper Cretaceous formations of southwestern Arkansas: Arkansas Geol. Survey Bull. 1, 213 p.
5. Danilchik, Walter, and Haley, B.R., 1964, Geology of the Paleozoic area in the Malvern quadrangle, Garland and Hot Spring Counties, Arkansas: U.S. Geol. Survey Misc. Geol. Inv. Map I–405.
6. Glick, E.E., and Frazon, S.E., 1965, Geologic map of The Snowball quadrangle, Newton and Searcy Counties, Arkansas: U.S. Geol. Survey Geol. Quad. Map GQ–425.
7. _____, U.S. Geol. Survey, unpublished data.
8. Glick, E.E., Haley, B.R., and Merewether, E.A., U.S. Geol. Survey, unpublished data.
9. Gordon, Mackenzie, Jr., and Kinney, D.M., 1964, Geologic map and structure sections of the Batesville district, Independence County, Arkansas: U.S.Geol. Survey Oil and Gas Inv. Prelim. Map 12.
10. Gordon, Mackenzie, Jr., Tinney, J.I., Jr., and Ellis, M.W., 1958, Geology of the Arkansas bauxite region: U.S. Geol. Survey Prof. Paper 299, 268 p.
11. Haley, B.R., 1961, Geology of Paris quadrangle, Logan County, Arkansas: Arkansas Geol. and Conser. Comm. Inf. Circ. 20–B, 40 p.
12. _____, 1966, Geology of the Barber quadrangle, Sebastian County and vicinity, Arkansas: Arkansas Geol. Comm. Inf. 20–C, 76 p.
13. _____, 1968, Geology of the Scranton and New Blaine quadrangles, Logan and Johnson Counties, Arkansas: U.S. Geol. Survey Prof. Paper 536–B, 10 p.
14. _____, U.S. Geol. Survey, unpublished data.
15. Haley, B.R., and Hendricks, T.A., 1968, Geology of the Greenwood quadrangle, Arkansas—Oklahoma: U.S.Geol. Survey Prof. Paper 536–A, 15 p.
16. _____1971, Geology of the Van Buren and Lavaca quadrangles, Arkansas and Oklahoma: U.S. Geol. Survey Prof. Paper 657–A, 41 p.
17. Haley, B.R., and Merewether, E.A., U.S. Geol. Survey, unpublished data.
18. Holbrook, D.F., Stone, C.G., and Sterling, P.J., Arkansas Geol. Comm., unpublished data.
19. Kinney, D.M., and Stracek, J.A., 1950, Geology map of the central part of the Batesville manganese district, Independence and Izard Counties, Arkansas: U.S. Geol. Survey Mineral Inv. Field Studies Map MF–1.
20. Maher, J.C., and Lantz, R.J., 1953, Geology of the Gilbert area, Searcy County, Arkansas: U.S. Geol. Survey Oil and Gas Inv. Map OM–132.
21. McKnight, E.T., 1935, Zinc and lead deposits of northern Arkansas: U.S. Geol. Survey Bull. 853, 311 p.
22. Merewether, E.A., 1971, Geology of the Knoxville and Delaware quadrangles, Johnson and Logan Counties and vicinity, Arkansas: U.S. Geol. Survey Prof. Paper 657–B, 18 p.
23. Mesavage, E.A., and Haley, B.R., 1969, Geology map of the Coal Hill, Hartman, and Clarksville quadrangles, Johnson County and vicinity, Arkansas: U.S. Geol. Survey Prof. Paper 536–C, 27 p.
24. Miser, H.D., and Purdue, A.H., 1929, Geology of the De Queen and Caddo Gap quadrangles, Arkansas: U.S. Geol. Survey Bull. 808, 195 p.
25. Purdue, A.H., 1907, Description of the Winslow quadrangle [Arkansas—Indian Territory]: U.S. Geol. Survey Geol. Atlas, Folio 154.
26. Purdue, A.H., and Miser, H.D., 1916, Description of the Eureka Springs and Harrison quadrangles [Arkansas—Missouri]: U.S. Geol. Survey Geol. Atlas, Folio 202.
27. _____1923, Description of the Hot Springs district [Arkansas]: U.S. Geol. Survey Geol. Atlas, Folio 215.
28. Reinemund, J.A., and Danilchik, Walter, 1957, Preliminary geologic map of the Waldron quadrangle and adjacent areas, Scott County, Arkansas: U.S. Geol. Survey Oil and Gas Inv. Map OM–192.
29. Renfroe, C.A., Arkansas Geol. Comm., unpublished data.
30. Stone, C.G., and Sterling, P.J., Arkansas Geol. Comm., unpublished data.
31. Wilbert, L.J., Jr., 1953, The Jacksonian Stage in southeastern Arkansas: Arkansas Resources Devel. Comm., Division of Geology Bull. 19, 125 p.

CHANGES IN STRATIGRAPHIC NAMES

Lithologic boundaries and stratigraphic nomenclature, when they differ from the Geologic Map of Arkansas (Arkansas Geological Survey, 1929), are summarized as follows:

GULF COASTAL PLAIN

The contact between the Claiborne Group and the Jacksonian Stage was established by Wilbert (1953). The stratigraphic rank was changed from formation to group for the Midway, Wilcox, Claiborne, and Jackson Groups by Gordon, Tracey, and Ellis (1958).

OZARK REGION

The Jefferson City Dolomite was first identified in Arkansas by E. O. Ulrich (in Purdue and Miser, 1916). The Jefferson City Dolomite and the overlying Cotter cannot be separated on the basis of lithology and are shown as a single unit on this map.

The Cotter Dolomite, the Powell Dolomite, and the Everton Formation were mapped in Marion County by McKnight (1935). These three formations were mapped eastward from Marion county to the Mississippi Embayment for this map. The upper part of the Powell as shown on this map in southeastern Sharp, northern Lawrence, and southern Randolph Counties includes rocks that were divided into the Smithville Limestone and the overlying Black Rock Limestone on the Geologic Map of Arkansas (1929).

Two map units, the Cane Hill Member of the Hale Formation, and the Prairie Grove Member of the Hale Formation and the Bloyd Shale, have been recognized eastward from the Eureka Springs section, southwestern Washington County, described by Hembest (1953). The Cane Hill Member is considered Pennsylvanian in age in the western part of Arkansas but it spans the Mississippian—Pennsylvanian boundary in the area east of Newton County, where fossils have been collected and identified as being of latest Mississippian age by Gordon (1964).

ARKANSAS VALLEY AND OUACHITA MOUNTAINS REGIONS

The ages of two formations are changed on this map: the Collier Shale is changed from Cambrian to Early Ordovician because it contains conodonts of that age (Repetski and Ethington, 1973); the Crystal Mountain Sandstone is changed from Ordovician(?) to Early Ordovician.

The upper part of the Stanley Shale includes beds of chert and siliceous shale equivalent to those in Oklahoma in the Chickasaw Creek Formation of Harlton (1938). The Hot Springs Sandstone is reduced in stratigraphic rank on this map to a member of the Stanley Shale. It is overlain and underlain by unnamed members of the Stanley in most places.

The Jackfork Sandstone is of Pennsylvanian (Morrowan) age in Arkansas, according to Gordon and Stone (1973).

The Johns Valley Shale near Boles, Scott County, was mapped in detail by Reinemund and Danilchik (1957). Erratic boulders from pre-Mississippian age formations are recognizable in Scott and extreme western Yell Counties, but not elsewhere.

The base of the McAlester Formation in Arkansas was established by Collier (1907); it coincides with the top of the lower sandstone member of the Hartshorne Sandstone in Oklahoma. The contacts between the McAlester, Savanna, and Boggy Formations were traced from their type localities in Oklahoma into Arkansas by Hendricks and Parks (1950), and were later modified by Haley (1961).

REFERENCES

Arkansas Geological Survey, 1929, Geologic map of Arkansas.
Collier, A.J., 1907, The Arkansas coal field, with reports on the paleontology, by David White and G.H. Girty: U.S. Geol. Survey Bull. 326, 156 p.
Gordon, Mackenzie, Jr., 1964, Moorefield formation and Ruddell shale, Batesville district, Arkansas: Am. Assoc. Petroleum Geologists Bull., v. 28, no. 11, p. 1626–1634.
_____1964, Carboniferous cephalopods of Arkansas: U.S. Geol. Survey Prof. Paper 460, 322 p.
Gordon, Mackenzie, Jr., and Stone, C.G., 1973, Correlation of Carboniferous rocks of the Ouachita geosyncline with those of the adjacent shelf: Geol. Soc. America Abs. with Programs, v. 5, no. 3, p. 259.
Gordon, Mackenzie, Jr., Tracey, J.I., Jr., and Ellis, M.W., 1958, Geology of the Arkansas bauxite region: U.S. Geol. Survey Prof. Paper 299, 268 p.
Haley, B.R., 1961, Geology of Paris quadrangle, Logan County, Arkansas: Arkansas Geol. and Conser. Comm. Inf. Circ. 20–B, 40 p.
Harlton, B.H., 1938, Stratigraphy of the Bendian of the Oklahoma salient of the Ouachita Mountains: Am. Assoc. Petroleum Geologists Bull., v. 22, no. 7, p. 852–914.
Hembest, L.G., 1953, Morrow group and lower Atoka formation of Arkansas: Am. Assoc. Petroleum Geologists Bull., v. 37, no. 8, p. 1935–1953.
Hendricks, T.A., and Parks, Bryan, 1950, Geology of the Fort Smith district, Arkansas: U.S. Geol. Survey Prof. Paper 221–E, p. 67–94.
McKnight, E.T., 1935, Zinc and lead deposits of northern Arkansas: U.S. Geol. Survey Bull. 853, 311 p.
Purdue, A.H., and Miser, H.D., 1916, Description of the Eureka Springs and Harrison quadrangles [Arkansas—Missouri]: U.S. Geol. Survey Geol. Atlas, Folio 202.
Reinemund, J.A., and Danilchik, Walter, 1957, Preliminary geologic map of the Waldron quadrangle and adjacent areas, Scott County, Arkansas: U.S. Geol. Survey Oil and Gas Inv. Map OM–192.
Repetski, J.E., and Ethington, R.L., 1973, Conodonts from graptolite facies in the Ouachita Mountains, Arkansas and Oklahoma: Geol. Soc. America Abs. with Programs, v. 5, no. 3, p. 277.
Stone, C.G., 1963, Notes on the eastermost frontal Ouachita-southeast Arkansas Valley area, Arkansas: Fort Smith Geol. Soc. 2d Regional Field Conf. Guidebook, 44 p.
Wilbert, L.J., Jr., 1953, The Jacksonian Stage in southeastern Arkansas: Arkansas Resources Devel. Comm., Division of Geology Bull. 19, 125 p.

PREPARED COOPERATIVELY BY THE
ARKANSAS GEOLOGICAL COMMISSION
NORMAN F. WILLIAMS, DIRECTOR
AND THE
UNITED STATES GEOLOGICAL SURVEY
DALLAS A. PECK, DIRECTOR

GEOLOGIC MAP OF ARKANSAS

By

Boyd R. Haley, U.S. Geological Survey

Assisted by Ernest E. Glick, U.S. Geological Survey
and William V. Bush, Benjamin F. Clardy, Charles G. Stone,
Mac B. Woodward, and Doy L. Zachry, Arkansas Geological Commission

1993
(Revised from the 1976 edition)

SCALE 1:500 000

10 0 10 20 30 40 50 MILES
10 0 10 20 30 40 50 60 70 KILOMETERS

NATIONAL GEODETIC VERTICAL DATUM OF 1929

Cartography directed by James Gueen, U.S. Geological Survey
Cartography linework and lettering by Lawrence Brence
Color separations by Jennifer Fowler and Tony Gaeto

For sale by U.S. Geological Survey, Map Distribution
Box 25286, Federal Center, Denver, CO 80225, and
Arkansas Geological Commission
3815 West Roosevelt Road, Little Rock, AR 72204